

Bactericide, Immunomodulating, and Wound Healing Properties of Transgenic *Kalanchoe pinnata* Synergize with Antimicrobial Peptide Cecropin P1 In Vivo

Lebedeva A., Zakharchenko N., Trubnikova E., Medvedeva O., Kuznetsova T., Masgutova G., Zylkova M., Buryanov Y., Belous A.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Procedure of manufacturing *K. pinnata* water extracts containing cecropin P1 (CecP1) from the formerly described transgenic plants is established. It included incubation of leaves at +4°C for 7 days, mechanical homogenization of leaves using water as extraction solvent, and heating at +70°C for inactivating plant enzymes. Yield of CecP1 (after heating and sterilizing filtration) was 0.3% of total protein in the extract. The water extract of *K. pinnata* + CecP1 exhibits favorable effect on healing of wounds infected with *S. aureus* (equal to Cefazolin) and with a combination of *S. aureus* with *P. aeruginosa* (better than Cefazolin). Wild-type *K. pinnata* extract exhibited evident microbicide activity against *S. aureus* with *P. aeruginosa* but it was substantially strengthened in *K. pinnata* + CecP1 extract. *K. pinnata* extracts (both wild-type and transgenic) did not exhibit general toxicity and accelerated wound recovery. Due to immunomodulating activity, wild-type *K. pinnata* extract accelerated granulation of the wound bed and marginal epithelialization even better than *K. pinnata* + CecP1 extract. Immunomodulating and microbicide activity of *K. pinnata* synergizes with microbicide activity of CecP1 accelerating elimination of bacteria.

<http://dx.doi.org/10.1155/2017/4645701>
